

DISCLAIMER

The present version of the national guideline has been accepted by the President of the CPVO for its use in technical examinations carried out on behalf of the CPVO or for the take-over of reports serving as a basis for a CPVO decision.

Ligustrum ibota Siebold
Simplified standard protocol: SSP/LII/2.rev

Botanical taxon: Ligustrum Ibota Siebold
 Common Name (when known): Ligustrum
 Date of preparation of SSP: 14-09-2012
 Date of revision of SSP: 22-02-2021
 SSP revised by: W.A. Wietsma

Sample to be examined: VEGETATIVE
 Number of growing cycles: 1 year

Closing date for applications: 1/12
 Submission date/period: 1/4 - 30/4
 Seed/Plant Quantity: 10 young plants of commercial standard
 Able to show all their characteristics in the first year of examination

Special conditions sample: None

Test station address: Naktuinbouw, Sotaweg 22, 2371 AA, Roelofarendsveen
 Name: Team Support Variety Testing Department
 E-mail: teamsupport@rasraad.nl

List of grouping characteristics: NO, (if yes put as annex)
 Minimum number of plants in trial: Vegetative: 8 Seed: not appl.
 Minimum number of plants observed by measuring or counting: Vegetative: 1 Seed: not appl.

Give description of when observations on the flower should take place:
 At full flowering (if present)
 Give description of when/where observations on the leaf should take place:
 End summer - autumn
 Give description of when/where the other observations should take place:
 End summer – autumn

Test will take place: OUTDOORS

Uniformity: A population standard of 1% with an acceptance probability of at least 95%. Number of Off-types allowed: one off-type allowed in a sample size of 10

Table of characteristics: PRESENT (see annex)
 (if present, please annex the table of characteristics and explanations)

Literature: PRESENT
 (when present, please annex to this document)

Table of characteristics Ligustrum

| |
|---|
| Plant: attitude creeping/upright |
| Plant: height |
| |
| Stem: consistence herbaceous/lignified |
| Stem: ramification absent/present |
| Stem: thickness at base |
| Stem: color |
| Stem: pubescence |
| Internodes: length |
| |
| Leaf: arrangement |
| Leaf: petiole: length |
| Leaf: Blade: length |
| Leaf: Blade: width |
| Leaf: Blade: shape |
| Leaf: Blade: color of upper side |
| Leaf: Blade: color of lower side |
| Leaf: Blade: shape in cross section (concave/flat/convex) |
| Leaf: Blade: shape between veins (plicate/flat) |
| Leaf: Blade: margin: type of incisions |
| Leaf: Blade: shape of apex |
| Leaf: Blade: shape of base |
| Leaf: Blade: shape in cross section |
| |
| <i>Inflorescence: if present</i> |
| Inflorescence: type |
| Peduncle: length |
| Peduncle: width |
| Flower: Pedicel: length |
| Flower: Pedicel: pubescence |
| Flower: Pedicel: color |
| Flower: Pedicel: anthocyanin coloration |
| Flower: Stipulae: length |
| Flower: width |
| Flower: color |
| Flower: single/double |
| Flower: pubescence |
| Flower: Calyx: number of lobes |
| Flower: Calyx: shape of lobes |
| Flower: Calyx: length |
| Flower: Calyx: width: |
| Flower: Calyx: color |
| Flower: Calyx: anthocyanin coloration |
| Flower: Calyx: pubescence |
| |
| Flower: Corolla: shape |
| Flower: Corolla: length |
| Flower: Corolla: width |
| Flower: Corolla tube: color of base |
| Flower: Corolla tube: color of top |
| Flower: Corolla tube: pubescence on inner side |
| Flower: Corolla: number of lobes |
| Corolla lobes: length |
| Corolla lobes: width |
| Corolla lobes: shape |

| |
|--|
| Corolla lobes: color |
| Stamen: number |
| stamen: length |
| stamen: color |
| anther: color |
| style: length |
| style: color |
| stigma: color |
| |
| Literature: Atrium tuinplanten encyclopedie: C. Brickell Hortica Color Cyclopedia of Garden Flora in all Climates and Plants indoor: A.B. Graf |