

DISCLAIMER

The present version of the national guideline has been accepted by the President of the CPVO for its use in technical examinations carried out on behalf of the CPVO or for the take-over of reports serving as a basis for a CPVO decision.

Tradescantia
Simplified standard protocol: SSP/TDC/3

Examination office	Naktuinbouw	
Reference of the protocol	SSP/TDC/3	
Date of preparation of the protocol	01/09/2021	
Date of entry into force of the protocol	01/09/2021	
Botanical taxon:	Tradescantia L. Tradescantia cerinthoides kunth Tradescantia albiflora Tradescantia spathacea Sw. Tradescantia virginiana L.	
Common Name (when known):	Spider wort Tradescantia Wandering-Jew Spider wort Tradescantia virginiana L.	
Way of propagation of the plants to be examined	Self or cross pollinated seed propagated <input type="checkbox"/> Vegetatively propagated <input checked="" type="checkbox"/>	
Number of growing cycles:	1 <input checked="" type="checkbox"/> 2 <input type="checkbox"/> Other <input type="checkbox"/> specify Click or tap here to enter text.	
List of grouping characteristics	Yes <input checked="" type="checkbox"/> if yes put as annex No <input type="checkbox"/>	
Minimum number of plants in trial	Vegetative:20	Seed: -
Minimum number of plants observed by measuring or counting:	Vegetative:1	Seed: -
Give description of when observations should take place	Observation on the flower should take place: at full flowering Observation on the leaf should take	

	<p>place: after 6 month growing in the test facility.</p> <p>Other observations should take place: after 6 month growing in the test facility.</p>
<p>Uniformity:</p> <ul style="list-style-type: none"> - For the assessment of uniformity of vegetatively propagated, self-pollinated seed propagated varieties or F1-hybrids, a population standard of 1 % and an acceptance probability of at least 95% should be applied. In the case of a sample size of 20 plants, 1 off-types are allowed. - For the assessment of uniformity for cross-pollinated varieties, the recommendations for cross-pollinated varieties in the General introduction of UPOV should be applied. The variability within the variety should not exceed the variability of comparable varieties already known. 	
<p>Table of characteristics</p>	<p>Present <input checked="" type="checkbox"/></p> <p>Not available <input type="checkbox"/></p>
<p>Literature (when present, please annex to this document)</p>	<p>Present <input checked="" type="checkbox"/></p> <p>Absent <input type="checkbox"/></p>

Table of characteristics: Tradescantia

1. Plant: height
2. Plant: diameter
3. Stem: length of internodes
4. Stem: thickness of internodes
5. Stem: colour
6. Stem: anthocyanin coloration
7. Sheath: length
8. Sheath: width
9. Sheath: colour
10. Sheath: anthocyanin coloration
11. Leaf blade: length
12. Leaf blade: width
13. Leaf blade: shape
14. Leaf blade: position of broadest part
15. Leaf blade: number of colours
16. Leaf blade: main colour of upper side
17. Leaf blade: secondary colour of upper side
18. Leaf blade: main colour of lower side
19. Leaf blade: pubescence
20. Leaf blade: shape of apex
21. Leaf blade: undulation of margin
22. Peduncle: length
23. Peduncle: colour
24. Calyx: length
25. Calyx: width
26. Calyx: colour
27. Calyx: anthocyanin coloration
28. Calyx: pubescence
29. Flower: diameter
30. Petal: length
31. Petal: width
32. Petal: shape of apex
33. Peta: number of colours
34. Petal: main colour RHS Colour Chart (indicate reference number)
35. Petal: secondary colour RHS Colour Chart (indicate reference number)
36. Anther: colour
37. Style: colour
Literature: Name that flower: I. Clarke, H. Lee Temperate Garden Plant Families: P. Goldblatt, J.C. Manning