

DISCLAIMER

The present version of the national guideline has been accepted by the President of the CPVO for its use in technical examinations carried out on behalf of the CPVO or for the take-over of reports serving as a basis for a CPVO decision.

Ficus
Simplified standard protocol: SSP/VGB/3.rev

Examination office:	Naktuinbouw
Reference of the protocol:	SSP/VGB/3.rev
Date of preparation of the protocol:	01/02/2023
Date of entry into force of the protocol:	01/03/2022
Botanical taxon:	<p>Ficus americana Aubl. subsp. guianensis (Ham.) C. C. Berg</p> <p>Ficus auriculata Lour.</p> <p>Ficus benghalensis L.</p> <p>Ficus binnendijkii Miq.</p> <p>Ficus bussei Warb. ex Mildbr. & Burret</p> <p>Ficus cyathistipula Warb.</p> <p>Ficus deltoidea Jack</p> <p>Ficus elastica Roxb. ex Hornem.</p> <p>Ficus L.</p> <p>Ficus lyrata Warb.</p> <p>Ficus microcarpa L. f.</p> <p>Ficus natalensis Hochst.</p> <p>Ficus petiolaris Kunth</p> <p>Ficus pumila L.</p> <p>Ficus punctata Thunb.</p> <p>Ficus religiosa L.</p> <p>Ficus sagittata Vahl</p> <p>Ficus stricta (Miq.) Miq.</p> <p>Ficus umbellata Vahl</p>
Common Name (when known):	<p>Ficus americana Aubl. subsp. Guianensis</p> <p>Roxburgh fig</p> <p>Ficus benghalensis</p> <p>Fig</p> <p>Ficus bussei</p> <p>Ficus cyathistipula</p>

	Ficus deltoidei Rubber Plant Fig tree Fiddle-leaf fig Ficus macrocarpa Bark-cloth fig Ficus petiolaris Fig pumila Ficus punctata Fig Ficus sagittate Ficus stricta Ficus umbellata	
Way of propagation of the plants to be examined:	Self or cross pollinated seed propagated <input type="checkbox"/> Vegetatively propagated <input checked="" type="checkbox"/>	
Number of growing cycles:	1 <input checked="" type="checkbox"/> 2 <input type="checkbox"/> Other <input type="checkbox"/> specify Click or tap here to enter text.	
List of grouping characteristics:	Yes <input type="checkbox"/> if yes put as annex No <input checked="" type="checkbox"/>	
Minimum number of plants in trial:	Vegetative:20	Seed: -
Minimum number of plants observed by measuring or counting:	Vegetative:1	Seed: -
Give description of when observations should take place:	Observation should take place: after 9 months of growing in the test facility	

<p>Uniformity:</p> <ul style="list-style-type: none"> - For the assessment of uniformity of vegetatively propagated, self-pollinated seed propagated varieties or F1-hybrids, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 20 plants, 1 off-type is allowed - For the assessment of uniformity for cross-pollinated varieties, the recommendations for cross-pollinated varieties in the General introduction of UPOV should be applied. The variability within the variety should not exceed the variability of comparable varieties already known. 	
Table of characteristics:	<p>Present <input checked="" type="checkbox"/></p> <p>Not available <input type="checkbox"/></p>
Literature: (when present, please annex to this document)	<p>Present <input type="checkbox"/></p> <p>Absent <input checked="" type="checkbox"/></p>

